

Calculation of Ultraviolet Radiation Quantities using TOMS Ozone and ISCCP Cloud Data

Bernhard Mayer and Sasha Madronich

NCAR, P.O. Box 3000, Boulder, CO 80307; e-mail bmayer@ucar.edu, sasha@ucar.edu

**National
Center for
Atmospheric
Research**

Acknowledgments

The authors would like to thank Chris Fischer for the TOMS-based calculations for Figure 4. B.M. was supported by a research grant from the German Academic Exchange Service (DAAD). S.M. was supported by the Department of Energy through grant No. DE-A105-94ER61879. NCAR is sponsored by the National Science Foundation. The maps were created with the free GMT package by P. Wessel and W. H. F. Smith.

1 Abstract

UV irradiance and actinic flux are calculated using two different approaches; (A) a well established TOMS cloud reflectivity method [1]; and (B) a new method, based on total ozone from TOMS and cloud parameters from ISCCP. Both products show good agreement between each other, and with ground-based measurements.

2 Methods

A. TOMS-based method. Surface radiation is calculated by scaling the cloudless sky value with a wavelength-independent factor γ which is derived from the TOMS reflectivity according to [1], [3]:

$$E = \gamma \cdot E_{\text{clear}}$$

B. ISCCP-based method. Radiation is calculated using cloud fraction and cloud optical depth from the ISCCP D1 data set [5]. Broken clouds are considered with the independent pixel approximation [4], [3]:


$$E = (1 - c) \cdot E_{\text{clear}} + c \cdot E_{\text{cloudy}}$$

where c is the cloud fraction, and E_{clear} and E_{cloudy} are the irradiances for cloudless and totally cloud-covered sky, calculated using the TUV radiative transfer model [2]. The snow cover information of the ISCCP D1 data is used to estimate the correct surface albedo. In both methods, cloudless sky irradiance is calculated using TOMS total ozone.



3 Results

Fig. 1 shows good agreement in the erythema irradiance calculated using both methods. While the TOMS-based method provides a quick way to calculate surface irradiance and even actinic flux or j -values (Fig. 2) [3], the ISCCP-based method might be of advantage for certain applications because: (1) some extra uncertainty is introduced when using TOMS reflectivity to calculate the actinic flux [3];

Erythema (CIE), 06/04/90, cloudy sky
TOMS



ISCCP




Irradiance [W / m^2]

Figure 1: TOMS– and ISCCP–based noontime erythemal irradiance.

$j[O(^1D)]$, 06/01/90, cloudy sky

TOMS



ISCCP





Figure 2: TOMS– and ISCCP–based noontime photolysis frequency $j[O(^1D)]$.

(2) the ISCCP data distinguish between clouds and surface albedo; (3) the ISCCP data include information about the vertical layering of the clouds; and (4) the time resolution of the IS–CCP data is higher.

The ISCCP D1 data are available in steps of 3 hours, thus allowing the calculation of diurnal variations. Here, nine data points, equally spaced in time and centered around local noon, are sufficient to calculate daily doses with an accuracy of 1% compared to a calculation with high time resolution. Fig. 3 shows a pronounced diurnal variation of cloudiness, which can only be obtained from the ISCCP data.

A comparison of the TOMS-based data with ground-based measurements from the World Ozone and UV data center (WOUDC) shows good agreement (Fig. 4). A systematic difference of 5–10% is expected because most of the measured data were not corrected for the cosine error of the entrance optics.




Figure 3: Diurnal variation.




Figure 4: Model calculations and measurements of averaged daily erythemal irradiation.

References

- [1] Eck, T., P. Barthia, and J. Kerr, *Geophys. Res. Lett.*, 22, 611–614, 1995.
- [2] Madronich, S., Environmental Effects of UV (Ultraviolet) Radiation, in *UV radiation in the natural and perturbed atmosphere*, Lewis Publisher, Boca Raton, 17–69, 1993.
- [3] Mayer, B., C.A. Fischer, and S. Madronich, *Geophys. Res. Lett.*, 25, 4321–4324, 1998.
- [4] Nack, M., and A. Green, *Appl. Opt.*, 13, 2405–2415, 1974.
- [5] Rossow, W. et al., *International Satellite Cloud Climatology Project (ISCCP) –Documentation of new cloud datasets*, <http://isccp.giss.nasa.gov>, 1996.